American Society of Mammalogists

Germination of Cow Parsnip Seeds from Grizzly Bear Feces Author(s): Roger D. Applegate, Lynn L. Rogers, David A. Casteel, James M. Novak Source: *Journal of Mammalogy*, Vol. 60, No. 3 (Aug., 1979), p. 655 Published by: American Society of Mammalogists Stable URL: <u>http://www.jstor.org/stable/1380120</u> Accessed: 16/01/2011 16:39

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at http://www.jstor.org/page/info/about/policies/terms.jsp. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at http://www.jstor.org/action/showPublisher?publisherCode=asm.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



American Society of Mammalogists is collaborating with JSTOR to digitize, preserve and extend access to Journal of Mammalogy.

GERMINATION OF COW PARSNIP SEEDS FROM GRIZZLY BEAR FECES

This is a report on the germinability of seeds retrieved from feces of grizzly bears (Ursus arctos). Impetus for this work was our observations of plants growing from grizzly feces on moraines in Glacier National Park and of seeds germinating in droppings of birds, raccoons (Procyon lotor), and oppossums (Didelphis virginiana). Additionally, Krefting and Roe (Ecol. Monogr., 19:270–286, 1949) noted that seeds of blueberries (Vaccinium spp.) sometimes germinated from bear feces and that seeds of wild sarsaparilla (Aralia nudicaulis) from black bear feces germinated better than controls. No tests of statistical significance were associated with the latter observation, however. The objective of this study was to determine if germination rates of seeds from bear feces were different from those of seeds from fruit picked when ripe. This information may help to evaluate the importance of bears as dispersers of plants.

Germination trials were conducted for 150 seeds of cow parsnip (*Heracleum lanatum*) collected from six grizzly bear fecal droppings in Glacier National Park, Montana, and for 150 seeds picked as controls. Half of the seeds from each group were frozen overwinter; the remainder were not frozen. All seeds were placed in a growth chamber in moist sand using temperature and light regimes simulating those of spring in Glacier National Park. Of the seeds from bear scats, 49 (65%) of those unfrozen and 64 (85%) of those frozen germinated. Of the control treatments, only 38 (51%) of the unfrozen and 52 (69%) of the frozen seeds germinated. Differences in germination rates for seeds in all treatments were significant at the 5% level using Chi-square. The higher rates of germination of cow parsnip seeds passed through the digestive tracts of bears, and the long distances traveled by grizzly bears (Craighead, Natl. Geogr. Soc. Res. Repts., 1963 projects:59–67, 1968), suggest that bears may be important dispersers of seeds.

ROGER D. APPLEGATE, LYNN L. ROGERS, DAVID A. CASTEEL, AND JAMES M. NOVAK, University of Illinois, Urbana, IL 61801, USDA Forest Service, North Central Forest Experiment Station, 1992 Folwell Avenue, St. Paul, MN 55108, USDI National Park Service, Glacier National Park, Babb, MT 59411, and Department of Forestry, University of Illinois, Urbana, IL 61801 (present address of Applegate: Tennessee Valley Authority, Division of Forestry, Norris, TN 37828). Submitted 15 November 1978. Accepted 29 December 1978.

J. Mamm., 60(3):655, 1979